

Lecture 16 – Equivalence of Pushdown Automata and Context-Free Grammars

COSE215: Theory of Computation

Jihyeok Park




2023 Spring

A **context-free grammar** is a 4-tuple:

$$G = (V, \Sigma, S, R)$$

A **pushdown automaton (PDA)** is a finite automaton with a **stack**.

- Acceptance by **final states**
- Acceptance by **empty stacks**



PDA_{FS}
(by final states)

|| ?

PDA_{ES}
(by empty stacks)

|| ?

CFG

1. Equivalence of PDA by Final States and Empty Stacks

PDA_{FS} to PDA_{ES}

PDA_{ES} to PDA_{FS}

2. Equivalence of PDA and CFGs

CFGs to PDA_{ES}

PDA_{ES} to CFGs



1. Equivalence of PDA by Final States and Empty Stacks


PDA_{FS} to PDA_{ES}

PDA_{ES} to PDA_{FS}

2. Equivalence of PDA and CFGs

CFGs to PDA_{ES}

PDA_{ES} to CFGs




Theorem (PDA_{FS} to PDA_{ES})

For a given PDA $P = (Q, \Sigma, \Gamma, \delta, q_0, Z, F)$, \exists PDA P' . $L_F(P) = L_E(P')$.

Theorem (PDA_{FS} to PDA_{ES})

For a given PDA $P = (Q, \Sigma, \Gamma, \delta, q_0, Z, F)$, \exists PDA P' . $L_F(P) = L_E(P')$.



Theorem (PDA_{FS} to PDA_{ES})

For a given PDA $P = (Q, \Sigma, \Gamma, \delta, q_0, Z, F)$, \exists PDA P' . $L_F(P) = L_E(P')$.


Define a PDA

$$P' = (Q \cup \{q'_0, q'_1\}, \Sigma, \Gamma \cup \{Z'\}, \delta', q'_0, Z', \emptyset)$$



where

$$\begin{aligned} \delta'(q'_0, \epsilon, Z') &= \{(q_0, ZZ')\} \\ \delta'(q \in Q, a \in \Sigma, X \in \Gamma) &= \delta(q, a, X) \\ \delta'(q \in Q, \epsilon, X \in \Gamma \cup \{Z'\}) &= \begin{cases} \delta(q, \epsilon, X) \cup \{(q'_1, \epsilon)\} & \text{if } q \in F \\ \delta(q, \epsilon, X) & \text{otherwise} \end{cases} \\ \delta'(q'_1, \epsilon, X \in \Gamma \cup \{Z'\}) &= \{(q'_1, \epsilon)\} \end{aligned}$$

$$L_F(P) = L_E(P') = \{a^n b^n \mid n \geq 0\}$$



$$L_F(P) = L_E(P') = \{a^n b^n \mid n \geq 0\}$$




Theorem (PDA_{ES} to PDA_{FS})

For a given PDA $P = (Q, \Sigma, \Gamma, \delta, q_0, Z, F)$, \exists PDA P' . $L_E(P) = L_F(P')$.

Theorem (PDA_{ES} to PDA_{FS})

For a given PDA $P = (Q, \Sigma, \Gamma, \delta, q_0, Z, F)$, \exists PDA P' . $L_E(P) = L_F(P')$.



Theorem (PDA_{ES} to PDA_{FS})

For a given PDA $P = (Q, \Sigma, \Gamma, \delta, q_0, Z, F)$, \exists PDA P' . $L_E(P) = L_F(P')$.

Define a PDA

$$P' = (Q \cup \{q'_0, q'_1\}, \Sigma, \Gamma \cup \{Z'\}, \delta', q'_0, Z', \{q'_1\})$$

where


$$\delta'(q'_0, \epsilon, Z') = \{(q_0, ZZ')\}$$

$$\delta'(q \in Q, a \in \Sigma, X \in \Gamma) = \delta(q, a, X)$$



$$\delta'(q \in Q, \epsilon, X \in \Gamma) = \delta(q, \epsilon, X)$$

$$\delta'(q \in Q, \epsilon, Z') = \{(q'_1, \epsilon)\}$$

$$L_E(P) = L_F(P') = \{a^n b^n \mid n \geq 0\}$$



$$L_E(P) = L_F(P') = \{a^n b^n \mid n \geq 0\}$$



1. Equivalence of PDA by Final States and Empty Stacks

PDA_{FS} to PDA_{ES}

PDA_{ES} to PDA_{FS}

2. Equivalence of PDA and CFGs

CFGs to PDA_{ES}

PDA_{ES} to CFGs



Theorem (CFGs to PDA_{ES})

For a given CFG $G = (V, \Sigma, S, R)$, \exists PDA P . $L(G) = L_E(P)$.

Define a PDA

$$P = (\{q\}, \Sigma, V \cup \Sigma, \delta, q, S, \emptyset)$$

where

$$\delta(q, \epsilon, A \in V) = \{(q, \alpha) \mid A \rightarrow \alpha \in R\}$$

$$\delta(q, a \in \Sigma, a \in \Sigma) = \{(q, \epsilon)\}$$

$$\begin{aligned}\delta(q, \epsilon, A \in V) &= \{(q, \alpha) \mid A \rightarrow \alpha \in R\} \\ \delta(q, a \in \Sigma, a \in \Sigma) &= \{(q, \epsilon)\}\end{aligned}$$

Consider the following CFG:

$$S \rightarrow \epsilon \mid aSb \mid bSa \mid SS$$


$$\begin{aligned} \delta(q, \epsilon, A \in V) &= \{(q, \alpha) \mid A \rightarrow \alpha \in R\} \\ \delta(q, a \in \Sigma, a \in \Sigma) &= \{(q, \epsilon)\} \end{aligned}$$

Consider the following CFG:

$$S \rightarrow \epsilon \mid aSb \mid bSa \mid SS$$

Then, the equivalent PDA (by empty stacks) is:

$\epsilon [S \rightarrow \epsilon]$
 $\epsilon [S \rightarrow aSb]$
 $\epsilon [S \rightarrow bSa]$
 $\epsilon [S \rightarrow SS]$
 $a [a \rightarrow \epsilon]$
 $b [b \rightarrow \epsilon]$




$$\begin{aligned} \delta(q, \epsilon, A \in V) &= \{(q, \alpha) \mid A \rightarrow \alpha \in R\} \\ \delta(q, a \in \Sigma, a \in \Sigma) &= \{(q, \epsilon)\} \end{aligned}$$

Consider the following CFG:

$$S \rightarrow \epsilon \mid aSb \mid bSa \mid SS$$

Then, the equivalent PDA (by empty stacks) is:

$\epsilon [S \rightarrow \epsilon]$	$(q, abab, S) \vdash (q, abab, aSb)$
$\epsilon [S \rightarrow aSb]$	$\vdash (q, bab, Sb)$
$\epsilon [S \rightarrow bSa]$	$\vdash (q, bab, bSab)$
$\epsilon [S \rightarrow SS]$	$\vdash (q, ab, Sab)$
$a [a \rightarrow \epsilon]$	$\vdash (q, ab, ab)$
$b [b \rightarrow \epsilon]$	$\vdash (q, b, b)$
	$\vdash (q, \epsilon, \epsilon)$



Theorem (PDA_{ES} to CFGs)

For a given PDA $P = (Q = \{q_0, \dots, q_{n-1}\}, \Sigma, \Gamma, \delta, q_0, Z, F)$,
 \exists CFG G . $L_E(P) = L(G)$.

Theorem (PDA_{ES} to CFGs)

For a given PDA $P = (Q = \{q_0, \dots, q_{n-1}\}, \Sigma, \Gamma, \delta, q_0, Z, F)$,
 \exists CFG G . $L_E(P) = L(G)$.

Consider the set of variables $V = \{S\} \cup \{A_{i,j}^X \mid 0 \leq i, j < n \wedge X \in \Gamma\}$.

Theorem (PDA_{ES} to CFGs)

For a given PDA $P = (Q = \{q_0, \dots, q_{n-1}\}, \Sigma, \Gamma, \delta, q_0, Z, F)$,
 \exists CFG G . $L_E(P) = L(G)$.

Consider the set of variables $V = \{S\} \cup \{A_{i,j}^X \mid 0 \leq i, j < n \wedge X \in \Gamma\}$.
 Then, define a CFG:

- For all $0 \leq j < n$,

$$S \rightarrow A_{0,j}^Z$$

- For all $q_i \in Q$, $a \in \Sigma \cup \{\epsilon\}$, and $X \in \Gamma$, consider any $(q_j, X_1 \cdots X_m) \in \delta(q_i, a, X)$ and $0 \leq k_1, \dots, k_m < n$. Then,

$$A_{i,k_m}^X \rightarrow a A_{j,k_1}^{X_1} A_{k_1,k_2}^{X_2} \cdots A_{k_{m-1},k_m}^{X_m}$$

Theorem (PDA_{ES} to CFGs)

For a given PDA $P = (Q = \{q_0, \dots, q_{n-1}\}, \Sigma, \Gamma, \delta, q_0, Z, F)$,
 \exists CFG G . $L_E(P) = L(G)$.

Consider the set of variables $V = \{S\} \cup \{A_{i,j}^X \mid 0 \leq i, j < n \wedge X \in \Gamma\}$.
 Then, define a CFG:

- For all $0 \leq j < n$,

$$S \rightarrow A_{0,j}^Z$$

- For all $q_i \in Q$, $a \in \Sigma \cup \{\epsilon\}$, and $X \in \Gamma$, consider any $(q_j, X_1 \cdots X_m) \in \delta(q_i, a, X)$ and $0 \leq k_1, \dots, k_m < n$. Then,


$$A_{i,k_m}^X \rightarrow a A_{j,k_1}^{X_1} A_{k_1,k_2}^{X_2} \cdots A_{k_{m-1},k_m}^{X_m}$$

Note that each variable $A_{i,j}^X$ generates all words that cause the PDA to go from state q_i to state q_j by popping X :

$$A_{i,j}^X \Rightarrow^* w \quad \text{if and only if} \quad (q_i, w, X) \vdash^* (q_j, \epsilon, \epsilon)$$


$$S \rightarrow A_{0,j}^Z \qquad A_{i,k_m}^X \rightarrow a A_{j,k_1}^{X_1} A_{k_1,k_2}^{X_2} \cdots A_{k_{m-1},k_m}^{X_m}$$

Consider the following PDA (by empty stacks):



$$S \rightarrow A_{0,j}^Z \qquad A_{i,k_m}^X \rightarrow a A_{j,k_1}^{X_1} A_{k_1,k_2}^{X_2} \cdots A_{k_{m-1},k_m}^{X_m}$$

Consider the following PDA (by empty stacks):




Then, the equivalent CFG is:

$$\begin{array}{l}
 S \rightarrow A_{0,0}^Z \mid A_{0,1}^Z \\
 A_{0,0}^Z \rightarrow a A_{0,0}^X A_{0,0}^Z \mid a A_{0,1}^X A_{1,0}^Z \mid A_{1,0}^Z \\
 A_{0,1}^Z \rightarrow a A_{0,0}^X A_{0,1}^Z \mid a A_{0,1}^X A_{1,1}^Z \mid A_{1,1}^Z \\
 A_{0,0}^X \rightarrow a A_{0,0}^X A_{0,0}^X \mid a A_{0,1}^X A_{1,0}^X \mid A_{1,0}^X \\
 A_{0,1}^X \rightarrow a A_{0,0}^X A_{0,1}^X \mid a A_{0,1}^X A_{1,1}^X \mid A_{1,1}^X \\
 A_{1,1}^Z \rightarrow \epsilon \\
 A_{1,1}^X \rightarrow b
 \end{array}$$

$$S \rightarrow A_{0,j}^Z \qquad A_{i,k_m}^X \rightarrow a A_{j,k_1}^{X_1} A_{k_1,k_2}^{X_2} \cdots A_{k_{m-1},k_m}^{X_m}$$

Consider the following PDA (by empty stacks):



Then, the equivalent CFG is:

$$\begin{array}{l}
 S \rightarrow A_{0,0}^Z \mid A_{0,1}^Z \\
 A_{0,0}^Z \rightarrow a A_{0,0}^X A_{0,0}^Z \mid a A_{0,1}^X A_{1,0}^Z \mid A_{1,0}^Z \\
 A_{0,1}^Z \rightarrow a A_{0,0}^X A_{0,1}^Z \mid a A_{0,1}^X A_{1,1}^Z \mid A_{1,1}^Z \\
 A_{0,0}^X \rightarrow a A_{0,0}^X A_{0,0}^X \mid a A_{0,1}^X A_{1,0}^X \mid A_{1,0}^X \\
 A_{0,1}^X \rightarrow a A_{0,0}^X A_{0,1}^X \mid a A_{0,1}^X A_{1,1}^X \mid A_{1,1}^X \\
 A_{1,1}^Z \rightarrow \epsilon \\
 A_{1,1}^X \rightarrow b
 \end{array}$$

$$\begin{array}{l}
 S \Rightarrow A_{0,1}^Z \\
 \Rightarrow a A_{0,1}^X A_{1,1}^Z \\
 \Rightarrow aa A_{0,1}^X A_{1,1}^X A_{1,1}^Z \\
 \Rightarrow aa A_{1,1}^X A_{1,1}^X A_{1,1}^Z \\
 \Rightarrow aab A_{1,1}^X A_{1,1}^Z \\
 \Rightarrow aabb A_{1,1}^Z \\
 \Rightarrow aabb
 \end{array}$$

1. Equivalence of PDA by Final States and Empty Stacks


PDA_{FS} to PDA_{ES}

PDA_{ES} to PDA_{FS}

2. Equivalence of PDA and CFGs

CFGs to PDA_{ES}

PDA_{ES} to CFGs



- Deterministic Pushdown Automata (DPDA)

Jihyeok Park

jihyeok_park@korea.ac.kr

<https://plrg.korea.ac.kr>