

# Lecture 9 – The Pumping Lemma for Regular Languages



COSE215: Theory of Computation

Jihyeok Park





2024 Spring

- Not all languages are regular: e.g.,  $L = \{a^n b^n \mid n \geq 0\}$ .





- Not all languages are regular: e.g.,  $L = \{a^n b^n \mid n \geq 0\}$ .



- How to prove that a language is **NOT** regular?

- Not all languages are regular: e.g.,  $L = \{a^n b^n \mid n \geq 0\}$ .



- How to prove that a language is **NOT** regular? **Pumping Lemma!**

## 1. Pumping Lemma for Regular Languages

Intuition

Pumping Lemma

Proof of Pumping Lemma

Application: Proving Languages are Not Regular

## 2. Examples

Example 1:  $L = \{a^n b^n \mid n \geq 0\}$

Example 2:  $L = \{ww^R \mid w \in \{a, b\}^*\}$

Example 3:  $L = \{a^l b^m c^n \mid l + m \leq n\}$

Example 4:  $L = \{a^{n^2} \mid n \geq 0\}$

Example 5:  $L = \{a^n b^k c^{n+k} \mid n, k \geq 0\}$

## 1. Pumping Lemma for Regular Languages

Intuition

Pumping Lemma

Proof of Pumping Lemma

Application: Proving Languages are Not Regular

## 2. Examples

Example 1:  $L = \{a^n b^n \mid n \geq 0\}$

Example 2:  $L = \{ww^R \mid w \in \{a, b\}^*\}$

Example 3:  $L = \{a^l b^m c^n \mid l + m \leq n\}$

Example 4:  $L = \{a^{n^2} \mid n \geq 0\}$

Example 5:  $L = \{a^n b^k c^{n+k} \mid n, k \geq 0\}$

# Intuition

Let's consider a **regular language**  $L$ .

Let's consider a **regular language**  $L$ .

Then, there exists a **regular expression**  $R$  such that  $L(R) = L$ .



Let's consider a **regular language**  $L$ .

Then, there exists a **regular expression**  $R$  such that  $L(R) = L$ .

Then, there are two possibilities:

Let's consider a **regular language**  $L$ .

Then, there exists a **regular expression**  $R$  such that  $L(R) = L$ .

Then, there are two possibilities:

- 1  $R$  does **not contain** the **Kleene star** operator ( $*$ ).

$abc$

$ab|(bc\epsilon)|d$

$(a|bcd)\epsilon|\emptyset bca$

Let's consider a **regular language**  $L$ .

Then, there exists a **regular expression**  $R$  such that  $L(R) = L$ .

Then, there are two possibilities:

- 1  $R$  does **not contain** the **Kleene star** operator ( $*$ ).

$abc$

$ab|(bc\epsilon)|d$

$(a|bcd)\epsilon|\emptyset bca$

Then, the language  $L$  should be **finite**.

Let's consider a **regular language**  $L$ .

Then, there exists a **regular expression**  $R$  such that  $L(R) = L$ .

Then, there are two possibilities:

- 1  $R$  does **not contain** the **Kleene star** operator ( $*$ ).

$abc$

$ab|(bc\epsilon)|d$

$(a|bcd)\epsilon|\emptyset bca$

Then, the language  $L$  should be **finite**.

- 2  $R$  **contains** at least one Kleene star operator ( $*$ ).

$$R_1 R_2^* R_3$$

Let's consider a **regular language**  $L$ .

Then, there exists a **regular expression**  $R$  such that  $L(R) = L$ .

Then, there are two possibilities:

- 1  $R$  does **not contain** the **Kleene star** operator ( $*$ ).

$abc$                        $ab|(bc\epsilon)|d$                        $(a|bcd)\epsilon|\emptyset bca$

Then, the language  $L$  should be **finite**.

- 2  $R$  **contains** at least one Kleene star operator ( $*$ ).

$$R_1 R_2^* R_3$$

Roughly speaking, we can **repeat** the middle part  $R_2$  as many times as we want (including 0 times).

Let's consider a **regular language**  $L$ .

Then, there exists a **regular expression**  $R$  such that  $L(R) = L$ .

Then, there are two possibilities:

- 1  $R$  does **not contain** the **Kleene star** operator ( $*$ ).

$abc \qquad ab|(bc\epsilon)|d \qquad (a|bcd)\epsilon|\emptyset bca$

Then, the language  $L$  should be **finite**.

- 2  $R$  **contains** at least one Kleene star operator ( $*$ ).

$$R_1 R_2^* R_3$$

Roughly speaking, we can **repeat** the middle part  $R_2$  as many times as we want (including 0 times).

The **Pumping Lemma** formally captures this intuition.

## Lemma (Pumping Lemma for Regular Languages)

For a given regular language  $L$ , **there exists** a *positive integer*  $n$  such that **for all**  $w \in L$ , if  $|w| \geq n$ , **there exists**  $w = xyz$  such that

- 1  $|y| > 0$
- 2  $|xy| \leq n$
- 3  $\forall i \geq 0. xy^iz \in L$

### Lemma (Pumping Lemma for Regular Languages)

For a given regular language  $L$ , **there exists** a *positive integer*  $n$  such that **for all**  $w \in L$ , if  $|w| \geq n$ , **there exists**  $w = xyz$  such that

- ①  $|y| > 0$
- ②  $|xy| \leq n$
- ③  $\forall i \geq 0. xy^iz \in L$

$A = L \text{ is regular}$



$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$



# Proof of Pumping Lemma

- Let  $L$  be a regular language.

# Proof of Pumping Lemma

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ .

# Proof of Pumping Lemma

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .

# Proof of Pumping Lemma

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ .

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ . Then,  $p_0 = q_0 \wedge p_m \in F$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ . Then,  $p_0 = q_0 \wedge p_m \in F$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\underbrace{\hspace{15em}}_{n+1 \text{ states}}$

- By Pigeonhole Principle, there exists  $i < j \leq n$  s.t.  $p_i = p_j$ .

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ . Then,  $p_0 = q_0 \wedge p_m \in F$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\underbrace{\hspace{15em}}_{n+1 \text{ states}}$

- By Pigeonhole Principle, there exists  $i < j \leq n$  s.t.  $p_i = p_j$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_i} p_i \xrightarrow{a_{i+1}} \cdots \xrightarrow{a_j} p_j \xrightarrow{a_{j+1}} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\overbrace{\hspace{10em}}^x \quad \overbrace{\hspace{10em}}^y \quad \overbrace{\hspace{15em}}^z$

- We can split  $w = xyz$  as above.



- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ . Then,  $p_0 = q_0 \wedge p_m \in F$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\underbrace{\hspace{15em}}_{n+1 \text{ states}}$

- By Pigeonhole Principle, there exists  $i < j \leq n$  s.t.  $p_i = p_j$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_i} p_i \xrightarrow{a_{i+1}} \cdots \xrightarrow{a_j} p_j \xrightarrow{a_{j+1}} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\overbrace{\hspace{10em}}^x$       $\overbrace{\hspace{10em}}^y$       $\overbrace{\hspace{15em}}^z$

- We can split  $w = xyz$  as above. Then,

$$|y| = j - i > 0 \qquad |xy| = j \leq n$$

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ . Then,  $p_0 = q_0 \wedge p_m \in F$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\underbrace{\hspace{15em}}_{n+1 \text{ states}}$

- By Pigeonhole Principle, there exists  $i < j \leq n$  s.t.  $p_i = p_j$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_i} p_i \xrightarrow{a_{i+1}} \cdots \xrightarrow{a_j} p_j \xrightarrow{a_{j+1}} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\overbrace{\hspace{10em}}^x \quad \overbrace{\hspace{10em}}^y \quad \overbrace{\hspace{15em}}^z$

- We can split  $w = xyz$  as above. Then,

$$|y| = j - i > 0 \qquad |xy| = j \leq n$$

- Since  $y$  represents a **cycle** from  $p_i$  to  $p_i$ ,  $\forall i \geq 0$ .  $\delta^*(q_0, xy^i z) = p_m$ .

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ . Then,  $p_0 = q_0 \wedge p_m \in F$ .

$$q_0 = \underbrace{p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} p_n}_{n+1 \text{ states}} \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

- By Pigeonhole Principle, there exists  $i < j \leq n$  s.t.  $p_i = p_j$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_i} \underbrace{p_i}_{x} \xrightarrow{a_{i+1}} \cdots \xrightarrow{a_j} \underbrace{p_j}_{y} \xrightarrow{a_{j+1}} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

- We can split  $w = xyz$  as above. Then,

$$|y| = j - i > 0 \qquad |xy| = j \leq n$$

- Since  $y$  represents a **cycle** from  $p_i$  to  $p_i$ ,  $\forall i \geq 0$ .  $\delta^*(q_0, xy^i z) = p_m$ .
- It means that  $\forall i \geq 0$ .  $xy^i z \in L$ .

- Let  $L$  be a regular language.
- Then,  $\exists$  DFA  $D = (Q, \Sigma, \delta, q_0, F)$ . s.t.  $L(D) = L$ . Let  $n = |Q| > 0$ .
- Take any  $w = a_1 a_2 \cdots a_m \in L$  s.t.  $|w| = m \geq n$ .
- Let  $p_i = \delta^*(q_0, a_1 \cdots a_i)$  for all  $0 \leq i \leq m$ . Then,  $p_0 = q_0 \wedge p_m \in F$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\underbrace{\hspace{15em}}_{n+1 \text{ states}}$

- By Pigeonhole Principle, there exists  $i < j \leq n$  s.t.  $p_i = p_j$ .

$$q_0 = p_0 \xrightarrow{a_1} p_1 \xrightarrow{a_2} \cdots \xrightarrow{a_i} p_i \xrightarrow{a_{i+1}} \cdots \xrightarrow{a_j} p_j \xrightarrow{a_{j+1}} \cdots \xrightarrow{a_n} p_n \xrightarrow{a_{n+1}} \cdots \xrightarrow{a_m} p_m \in F$$

$\overbrace{\hspace{10em}}^x \quad \overbrace{\hspace{10em}}^y \quad \overbrace{\hspace{15em}}^z$

- We can split  $w = xyz$  as above. Then,

$$\textcircled{1} - |y| = j - i > 0 \qquad |xy| = j \leq n - \textcircled{2}$$

- Since  $y$  represents a **cycle** from  $p_i$  to  $p_i$ ,  $\forall i \geq 0$ .  $\delta^*(q_0, xy^i z) = p_m$ .
- It means that  $\forall i \geq 0$ .  $xy^i z \in L$ . -  $\textcircled{3}$

## Lemma (Pumping Lemma for Regular Languages)

For a given regular language  $L$ , **there exists** a *positive integer*  $n$  such that **for all**  $w \in L$ , if  $|w| \geq n$ , **there exists**  $w = xyz$  such that

- ①  $|y| > 0$
- ②  $|xy| \leq n$
- ③  $\forall i \geq 0. xy^iz \in L$

$A = L \text{ is regular}$



$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$

## Lemma (Pumping Lemma for Regular Languages)

 $A = L \text{ is regular}$  $B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$

### Lemma (Pumping Lemma for Regular Languages)

$A = L \text{ is regular}$

$\Downarrow$

$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$

$A \implies B \quad (0)$

### Lemma (Pumping Lemma for Regular Languages)

$A = L \text{ is regular}$



$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$

$A \implies B \quad (O)$

$B \implies A \quad (X)$



### Lemma (Pumping Lemma for Regular Languages)

$A = L \text{ is regular}$

$\Downarrow$

$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$

$A \implies B \quad (0)$

$B \implies A \quad (X)$

$\neg B \implies \neg A \quad (0)$

## Lemma (Pumping Lemma for Regular Languages)

$$A = L \text{ is regular}$$



$$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$$

$$A \implies B \quad (0)$$

$$B \implies A \quad (X)$$

$$\neg B \implies \neg A \quad (0)$$

$$\neg B$$

### Lemma (Pumping Lemma for Regular Languages)

$$A = L \text{ is regular}$$



$$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$$

$$A \implies B \quad (0)$$

$$B \implies A \quad (X)$$

$$\neg B \implies \neg A \quad (0)$$

$$\begin{aligned} \neg B &= \neg(\exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \neg(\forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \end{aligned}$$

## Lemma (Pumping Lemma for Regular Languages)

$$A = L \text{ is regular}$$



$$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$$

$$A \implies B \quad (O)$$

$$B \implies A \quad (X)$$

$$\neg B \implies \neg A \quad (O)$$

$$\neg B = \neg(\exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3})$$

$$= \forall n > 0. \neg(\forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3})$$

$$= \forall n > 0. \exists w \in L. \neg(|w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3})$$

## Lemma (Pumping Lemma for Regular Languages)

$$A = L \text{ is regular}$$



$$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$$

$$A \implies B \quad (O)$$

$$B \implies A \quad (X)$$

$$\neg B \implies \neg A \quad (O)$$

$$\neg B = \neg(\exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3})$$

$$= \forall n > 0. \neg(\forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3})$$

$$= \forall n > 0. \exists w \in L. \neg(|w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3})$$

$$= \forall n > 0. \exists w \in L. |w| \geq n \wedge \neg(\exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3})$$

## Lemma (Pumping Lemma for Regular Languages)

$$A = L \text{ is regular}$$



$$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$$

$$A \implies B \quad (O)$$

$$B \implies A \quad (X)$$

$$\neg B \implies \neg A \quad (O)$$

$$\begin{aligned} \neg B &= \neg(\exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \neg(\forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. \neg(|w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \neg(\exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. \neg(\textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \end{aligned}$$

## Lemma (Pumping Lemma for Regular Languages)

$$A = L \text{ is regular}$$



$$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$$

$$A \implies B \quad (O)$$

$$B \implies A \quad (X)$$

$$\neg B \implies \neg A \quad (O)$$

$$\begin{aligned} \neg B &= \neg(\exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \neg(\forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. \neg(|w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \neg(\exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. \neg(\textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. \neg((\textcircled{1} \wedge \textcircled{2}) \vee \neg\textcircled{3}) \end{aligned}$$

## Lemma (Pumping Lemma for Regular Languages)

$$A = L \text{ is regular}$$



$$B = \exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}$$

$$A \implies B \quad (O)$$

$$B \implies A \quad (X)$$

$$\neg B \implies \neg A \quad (O)$$

$$\begin{aligned} \neg B &= \neg(\exists n > 0. \forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \neg(\forall w \in L. |w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. \neg(|w| \geq n \Rightarrow \exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \neg(\exists w = xyz. \textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. \neg(\textcircled{1} \wedge \textcircled{2} \wedge \textcircled{3}) \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. \neg(\textcircled{1} \wedge \textcircled{2}) \vee \neg\textcircled{3} \\ &= \forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. (\textcircled{1} \wedge \textcircled{2}) \Rightarrow \neg\textcircled{3} \end{aligned}$$



To prove a language  $L$  is **NOT** regular, we need to show that

$$\forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. (\textcircled{1} \wedge \textcircled{2}) \Rightarrow \neg \textcircled{3}$$

$$\textcircled{1} \quad |y| > 0$$

$$\textcircled{2} \quad |xy| \leq n$$

$$\textcircled{3} \quad \forall i \geq 0. xy^i z \in L$$

Note that  $\neg \textcircled{3} = \exists i \geq 0. xy^i z \notin L.$

To prove a language  $L$  is **NOT** regular, we need to show that

$$\forall n > 0. \exists w \in L. |w| \geq n \wedge \forall w = xyz. (\textcircled{1} \wedge \textcircled{2}) \Rightarrow \neg \textcircled{3}$$

- ①  $|y| > 0$
- ②  $|xy| \leq n$
- ③  $\forall i \geq 0. xy^i z \in L$

Note that  $\neg \textcircled{3} = \exists i \geq 0. xy^i z \notin L$ .

We can prove this by following the steps below:

- ① Assume **any** positive integer  $n$  is given.
- ② **Pick** a word  $w \in L$ .
- ③ Show that  $|w| \geq n$ .
- ④ Assume **any** split  $w = xyz$  is given, and  $\textcircled{1} |y| > 0 \wedge \textcircled{2} |xy| \leq n$ .
- ⑤  $\neg \textcircled{3}$  **Pick**  $i \geq 0$ , and show that  $xy^i z \notin L$  using  $\textcircled{1}$  and  $\textcircled{2}$ .

## 1. Pumping Lemma for Regular Languages

Intuition

Pumping Lemma

Proof of Pumping Lemma

Application: Proving Languages are Not Regular

## 2. Examples

Example 1:  $L = \{a^n b^n \mid n \geq 0\}$

Example 2:  $L = \{ww^R \mid w \in \{a, b\}^*\}$

Example 3:  $L = \{a^l b^m c^n \mid l + m \leq n\}$

Example 4:  $L = \{a^{n^2} \mid n \geq 0\}$

Example 5:  $L = \{a^n b^k c^{n+k} \mid n, k \geq 0\}$

## Example 1

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^n b^n \mid n \geq 0\}$$

## Example 1

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^n b^n \mid n \geq 0\}$$

- 1 Assume **any** positive integer  $n$  is given.

## Example 1

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^n b^n \mid n \geq 0\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^n b^n \in L$ .
- 3  $|w| = n + n = 2n \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .

# Example 1

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^n b^n \mid n \geq 0\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^n b^n \in L$ .
- 3  $|w| = n + n = 2n \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .
- 5 Let  $i = 0$ . We need to show that  $\neg$ ③  $xy^0z \notin L$ :

- Since ②  $|xy| \leq n$ ,

$$x = a^p \quad y = a^q \quad z = a^{n-p-q} b^n$$

for some  $0 \leq p, q \leq n$  such that  $p + q \leq n$ .

- Since ①  $|y| > 0$ , we know  $q > 0$ .
- Finally,  $xy^0z = xz = a^p a^{n-p-q} b^n = a^{n-q} b^n \notin L$  ( $\because q > 0$ ). □

## Example 2

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{ww^R \mid w \in \{a, b\}^*\}$$



## Example 2

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{ww^R \mid w \in \{a, b\}^*\}$$

- 1 Assume **any** positive integer  $n$  is given.

## Example 2

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{ww^R \mid w \in \{a, b\}^*\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^n b^n b^n a^n \in L$ .
- 3  $|w| = n + n + n + n = 4n \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .

## Example 2

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{ww^R \mid w \in \{a, b\}^*\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^n b^n b^n a^n \in L$ .
- 3  $|w| = n + n + n + n = 4n \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .
- 5 Let  $i = 0$ . We need to show that  $\neg$ ③  $xy^0z \notin L$ :
  - Since ②  $|xy| \leq n$ ,

$$x = a^p \quad y = a^q \quad z = a^{n-p-q} b^n b^n a^n$$

for some  $0 \leq p, q \leq n$  such that  $p + q \leq n$ .

- Since ①  $|y| > 0$ , we know  $q > 0$ .
- Finally,  $xy^0z = xz = a^p a^{n-p-q} b^n b^n a^n = a^{n-q} b^n b^n a^n \notin L$   
( $\because q > 0$ ).

□

## Example 3

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^l b^m c^n \mid l + m \leq n\}$$

## Example 3

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^l b^m c^n \mid l + m \leq n\}$$

- 1 Assume **any** positive integer  $n$  is given.

## Example 3

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^l b^m c^n \mid l + m \leq n\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^n b^n c^{2n} \in L$ .
- 3  $|w| = n + n + 2n = 4n \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .

## Example 3

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^l b^m c^n \mid l + m \leq n\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^n b^n c^{2n} \in L$ .
- 3  $|w| = n + n + 2n = 4n \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .
- 5 Let  $i = 2$ . We need to show that  $\neg$ ③  $xy^2z \notin L$ :
  - Since ②  $|xy| \leq n$ ,

$$x = a^p \quad y = a^q \quad z = a^{n-p-q} b^n c^{2n}$$

for some  $0 \leq p, q \leq n$  such that  $p + q \leq n$ .

- Since ①  $|y| > 0$ , we know  $q > 0$ .
- Finally,  $xy^2z = xyyz = a^{n+q} b^n c^{2n} \notin L$   
( $\because q > 0$ . Thus,  $(n + q) + n = 2n + q > 2n$ ).

□

## Example 4

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^{n^2} \mid n \geq 0\}$$



## Example 4

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^{n^2} \mid n \geq 0\}$$

- 1 Assume **any** positive integer  $n$  is given.

## Example 4

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^{n^2} \mid n \geq 0\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^{n^2} \in L$ .
- 3  $|w| = n^2 \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .

## Example 4

Let's prove that  $L$  is **NOT** regular using the Pumping Lemma:

$$L = \{a^{n^2} \mid n \geq 0\}$$

- 1 Assume any positive integer  $n$  is given.
- 2 Let  $w = a^{n^2} \in L$ .
- 3  $|w| = n^2 \geq n$ .
- 4 Assume any split  $w = xyz$  is given, and ①  $|y| > 0 \wedge$  ②  $|xy| \leq n$ .
- 5 Let  $i = 2$ . We need to show that  $\neg$ ③  $xy^2z \notin L$ :
  - Since ①  $|y| > 0$  and ②  $|xy| \leq n$ ,

$$y = a^k$$

where  $0 < k \leq n$ . Then,

$$n^2 < n^2 + k (\because 0 < k) \quad n^2 + k < (n + 1)^2 (\because k \leq n)$$

- Finally,  $xy^2z = xyyz = a^{n^2+k} \notin L$

□

## Example 5

Let's prove that  $L$  is **NOT** regular:

$$L = \{a^n b^k c^{n+k} \mid n, k \geq 0\}$$

## Example 5

Let's prove that  $L$  is **NOT** regular:

$$L = \{a^n b^k c^{n+k} \mid n, k \geq 0\}$$

- It is much easier to use **closure properties** under **homomorphisms**.
- Consider a homomorphism  $h : \{a, b, c\} \rightarrow \{a, b\}^*$ :

$$h(a) = a \quad h(b) = a \quad h(c) = b$$

- Then,

$$h(L) = \{a^{n+k} b^{n+k} \mid n, k \geq 0\} = \{a^n b^n \mid n \geq 0\}$$

- If  $L$  is regular, then  $h(L)$  must be regular as well.
- However, we know  $h(L)$  is **NOT** regular.
- Therefore,  $L$  is **NOT** regular. □

## 1. Pumping Lemma for Regular Languages

Intuition

Pumping Lemma

Proof of Pumping Lemma

Application: Proving Languages are Not Regular

## 2. Examples

Example 1:  $L = \{a^n b^n \mid n \geq 0\}$

Example 2:  $L = \{ww^R \mid w \in \{a, b\}^*\}$

Example 3:  $L = \{a^l b^m c^n \mid l + m \leq n\}$

Example 4:  $L = \{a^{n^2} \mid n \geq 0\}$

Example 5:  $L = \{a^n b^k c^{n+k} \mid n, k \geq 0\}$

- Please see this document on GitHub:

<https://github.com/ku-plrg-classroom/docs/tree/main/cose215/equiv-re-fa>

- The due date is 23:59 on Apr. 17 (Wed.).
- Please implement the following functions in `Implementation.scala`.
  - `reToENFA` for the conversion from REs to  $\epsilon$ -NFAs.
  - `dfaToRE` for the conversion from DFAs to REs.
  - `enfaToDFA` for the conversion from  $\epsilon$ -NFAs to DFAs.
- Please only submit `Implementation.scala` file to [Blackboard](#).

- Equivalence and Minimization of Finite Automata

Jihyeok Park

[jihyeok\\_park@korea.ac.kr](mailto:jihyeok_park@korea.ac.kr)

<https://plrg.korea.ac.kr>